

****FULL TITLE****

*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

Mass loss from red giants in the Magellanic Clouds

Jacco Th. van Loon

*Astrophysics Group, School of Physical and Geographical Sciences,
Keele University, Staffordshire ST5 5BG, UK*

Abstract. The Magellanic Clouds provide the ideal laboratory for the study of mass loss and molecule and dust formation in red giants as a function of luminosity and metal abundance. I present some recent observational work.

1. Mass-loss rates and wind kinematics

Mid-infrared surveys (IRAS, MSX) have resulted in the selection of hundreds of circumstellar dust envelopes in the Magellanic Clouds. Subsequent optical and infrared spectroscopy and photometry are used to identify the underlying stars, most of which are Asymptotic Giant Branch (AGB) stars or red supergiants (van Loon et al. 1997, 1998a; Trams et al. 1999b) that exhibit strong radial pulsations (Whitelock et al. 2003). The mass-loss rates are determined by modelling the spectral energy distributions, and found to depend on luminosity and stellar effective temperature (van Loon et al. 1999, 2005) as predicted by dust-driven wind theory including multiple scattering and in accordance with the results of hydrodynamical models. That theory also correctly predicts the wind speed as measured from circumstellar maser emission (van Loon et al. 1996, 1998b, 2001b; Marshall et al. 2004). Although the dust-to-gas ratio is proportional to metal abundance the total mass-loss rate may not depend sensitively on the initial metal abundance (van Loon 2000). The dependence of the mass-loss rate on initial mass and metal abundance can be calibrated by observation of mass-losing red giants that are members of star clusters within the Magellanic Clouds (van Loon et al. 2001a, 2003; van Loon et al. 2005a, in preparation).

2. Dust mineralogy and molecule formation

Surprisingly, both oxygen-rich AGB stars and red supergiants and carbon stars are found in the Magellanic Clouds that have very cool atmospheres despite their low initial metal abundance (van Loon et al. 1998a, 2005). In particular, it was noted that the carbonaceous molecular absorption bands in the $3\mu\text{m}$ spectra of magellanic carbon stars are at least as strong as in comparable galactic carbon stars (van Loon, Zijlstra & Groenewegen 1999; Matsuura et al. 2002, 2005). This may be due to the fact that carbon stars produce most of the carbon themselves. Massive oxygen-rich AGB stars do not become carbon stars due to Hot Bottom Burning (van Loon et al. 2001a: $M > 4M_{\odot}$; cf. Trams et al. 1999a), but they produce oxygen that may also reach the stellar surface and sustain the formation of copious amounts of molecules. However, apparently the large

molecular column density is not reflected in an equally high efficiency of dust grain condensation (see above), and there appears to be no strong connection between the strength of the dust continuum emission and the strength of the molecular absorption bands (van Loon et al. 2005b, in preparation). The dust detected in AGB stars and red supergiants in the LMC is similar to that in galactic sources (Trams et al. 1999b), with amorphous silicates dominating the mid-infrared spectrum of oxygen-rich sources and silicon-carbide seen in carbon stars, but detailed spectroscopic observations with the Spitzer Space Telescope (e.g., programme #3505 — PI: Peter Wood) are expected to reveal more subtle differences especially beyond $14\,\mu\text{m}$.




Figure 1. Average 2.9–4.1 μm VLT spectrum, for a sample of heavily dust-enshrouded carbon stars in the LMC (van Loon et al. 2005b, in preparation). Where the optical spectrum is dominated by diatomic molecules such as C_2 and CN , here we see more complex molecules including C_2H_2 and HCN .

References

- Marshall J.R., van Loon J.Th., Matsuura M., et al. 2004, MNRAS 355, 1348
Matsuura M., Zijlstra A.A., van Loon J.Th., et al. 2005, A&A in press
Matsuura M., Zijlstra A.A., van Loon J.Th., et al. 2002, ApJ 580, L133
Trams N.R., van Loon J.Th., Zijlstra A.A., et al. 1999a, A&A 344, L17
Trams N.R., van Loon J.Th., Waters L.B.F.M., et al. 1999b, A&A 346, 843
van Loon J.Th. 2000, A&A 354, 125
van Loon J.Th., Zijlstra A.A., Groenewegen M.A.T. 1999, A&A 346, 805
van Loon J.Th., Zijlstra A.A., Bujarrabal V., Nyman L.-Å 1996, A&A 306, L29
van Loon J.Th., Zijlstra A.A., Whitelock P.A., et al. 1997, A&A 325, 585
van Loon J.Th., Zijlstra A.A., Whitelock P.A., et al. 1998a, A&A 329, 169
van Loon J.Th., te Lintel Hekkert P., Bujarrabal V., et al. 1998b, A&A 337, 141
van Loon J.Th., Groenewegen M.A.T., de Koter A., et al. 1999, A&A 351, 559
van Loon J.Th., Zijlstra A.A., Kaper L., et al. 2001a, A&A 368, 239
van Loon J.Th., Zijlstra A.A., Bujarrabal V., Nyman L.-Å 2001b, A&A 368, 950
van Loon J.Th., Marshall J.R., Matsuura M., Zijlstra A.A. 2003, MNRAS 341, 1205
van Loon J.Th., Cioni M.-R.L., Zijlstra A.A., Loup C. 2005, A&A submitted
Whitelock P.A., Feast M.W., van Loon J.Th., Zijlstra A.A. 2003, MNRAS 342, 86